

Critical coupling in (1+1)-dimensional light-front ϕ^4 theory

Kazuto Oshima* and Masanobu Yahiro†

Gunma National College of Technology, Maebashi 371-8530, Japan

February 1, 2008

Abstract

The spontaneous symmetry breaking in (1+1)-dimensional ϕ^4 theory is studied with discretized light-front quantization, that is, by solving the zero-mode constraint equation. The symmetric ordering is assumed for the operator-valued constraint equation. The commutation relation between the zero mode and each oscillator mode is calculated with $\hbar$ expansion. A critical coupling evaluated from the first some terms in the expansion is $28.8\mu^2/\hbar \leq \lambda_{cr} \leq 31.1\mu^2/\hbar$ consistent with the equal-time one $22\mu^2/\hbar \leq \lambda_{cr} \leq 55.5\mu^2/\hbar$. The same analysis is also made under another operator ordering.

PACS numbers: 11.10.Lm, 11.30.Qc

*E-mail: oshima@nat.gunma-ct.ac.jp

†E-mail: yahiro@nat.gunma-ct.ac.jp

The light-front formulation of field theory is formally equivalent to the equal-time formulation within the framework of perturbation theory [1]. Nonperturbative calculations in the light-front formulation are successful in lower-dimensional field theories and yield the correct particle spectra [2, 3, 4]. The light-front formulation is then expected to be a promising method for solving the nonperturbative dynamics such as QCD. Nevertheless, the application to spontaneous symmetry breaking (SSB) is obstructed by the unsolved problem of the vacuum structure on the light cone [5, 6, 7]. It is then an important issue to confirm that for SSB the light-front formulation yields the same result as the equal-time formulation.

We study spontaneous breaking of the Z_2 symmetry in (1+1)-dimensional ϕ^4 theory as an instance. It is well known in the equal-time formulation that the model undergoes a phase transition at strong coupling [8]. In light-front field theory, the true vacuum always equals to the Fock vacuum [5], so SSB is believed to occur through the zero mode; a non-zero vacuum expectation value of the zero mode signals SSB. To ascertain this idea, several authors [9, 10, 11, 12, 13] have computed a vacuum expectation value of the zero mode and a critical coupling with either a truncation of Fock space or perturbation. Their results are qualitatively satisfactory. However, it is not clear whether the prescriptions are reliable for nonperturbative phenomena such as SSB, and it is not easy to improve the accuracy of the calculations. Our purpose in this brief report is to propose a different computation rule and obtain more accurate results. This rule yields a systematic way of obtaining a critical coupling λ_{cr} and a critical exponent β .

We define the light-front coordinates $x^\pm = (x^0 \pm x^1)/\sqrt{2}$. The Lagrangian density of (1 + 1)-dimensional ϕ^4 theory is described as

$$\mathcal{L} = \partial_+ \phi \partial_- \phi - \frac{\mu^2}{2} \phi^2 - \frac{\lambda}{4!} \phi^4. \quad (1)$$

We put the quantum system in a box of length d and impose periodic boundary conditions.

The field operator ϕ is then expanded as

$$\phi(x) = \frac{1}{\sqrt{4\pi}}a_0 + \sum_{n \neq 0} \frac{1}{\sqrt{4\pi|n|}}a_n(x^+)e^{ik_n^+x^-}, \quad (2)$$

where $k_n^+ = 2\pi n/d$. Coefficients of the expansion are operators which satisfy the canonical commutation relations [5, 11, 12],

$$[a_k, a_l] = [a_k^\dagger, a_l^\dagger] = 0, \quad [a_k, a_l^\dagger] = \hbar\delta_{k,l}, \quad k, l > 0, \quad (3)$$

where $a_k^\dagger = a_{-k}$. The zero-mode operator a_0 is not an independent quantity, since it governs the constraint equation [5, 11]

$$\begin{aligned} 0 = & a_0^3 + ga_0 + \sum_{n \geq 1} \frac{1}{n} (a_0 a_n a_n^\dagger + a_0 a_n^\dagger a_n + a_n a_n^\dagger a_0 + a_n^\dagger a_n a_0 + a_n a_0 a_n^\dagger + a_n^\dagger a_0 a_n \\ & - 3\hbar a_0) + \sum_{k,l,m \neq 0} \frac{\delta_{k+l+m,0}}{\sqrt{|klm|}} a_k a_l a_m \equiv \Phi(a, a^\dagger), \end{aligned} \quad (4)$$

where we have added the term $\sum_{n \geq 1} \frac{-3}{n}\hbar a_0$ to remove the tadpole divergences. The commutation relations (3) and the constraint equation (4) are obtained by quantizing the classical system with the Dirac-Bergmann quantization procedure. The procedure does not specify an operator ordering in the constraint equation (4), so we have assumed a Weyl (symmetric) ordering [14]. This assumption is reasonable, since it describes the phase transition properly, as shown later. The zero mode a_0 is a complicated operator obeying Eq.(4). We see from the quantization procedure that the zero mode satisfies [11]

$$[a_0, a_m^\dagger a_m] = 0. \quad (5)$$

The commutation relation guarantees that a_0 preserves the longitudinal momentum conservation.

So far the operator relation $[a_0, a_n]$ has not been esteemed significant. In literature, the corresponding classical relation (Dirac bracket) is derived from a classical constraint equation corresponding to Eq.(4), but it is impossible to quantize the Dirac bracket by simply replacing it by the commutation relation. Because, the quantization procedure

does not specify an operator ordering for operator products appearing in $[a_0, a_n]$. However, the importance of $[a_0, a_n]$ should be emphasized. In fact, as soon as $[a_0, a_n]$ is obtained, we can calculate $\sigma \equiv \langle 0|a_0|0\rangle$ as a function of $g \equiv \frac{24\pi\mu^2}{\lambda}$, as shown later. We derive the $[a_0, a_n]$ with Eq.(5) from

$$[\Phi(a, a^\dagger), a_n] = 0, \quad (6)$$

expanding $[a_0, a_n]$ as

$$[a_0, a_n] \equiv \sum_{p \geq 1} \tilde{\alpha}_p(n) \hbar^p. \quad (7)$$

If $[a_0, a_n]$ is obtained without the expansion, it should be a function of $a_0, a_n, a_n^\dagger$ and $\hbar$. Expanding the operator-valued function, we see that the expansion series is not unique because of the ambiguity on operator ordering. We only obtain $\tilde{\alpha}_p$ uniquely, when we take an operator ordering for $\tilde{\alpha}_p$. Since there would be no a priori proper operator-ordering for $\tilde{\alpha}_p$, we take a simple operator ordering shown below. Difference in operator ordering for $\tilde{\alpha}_p$ affects only to the higher order terms $\tilde{\alpha}_q$ ($q > p$).

At $O(\hbar)$, we have $[a_0^3, a_n] = 3a_0^2\tilde{\alpha}_1(n)\hbar + O(\hbar^2)$ etc., and from Eq.(6) we obtain

$$\tilde{\alpha}_1(n) = \frac{6}{nA}a_0a_n + \frac{3}{A} \sum_{k \neq 0, n} \frac{1}{\sqrt{|kn(n-k)|}} a_k a_{n-k}, \quad (8)$$

where

$$A = g + 3a_0^2 + \sum_{m \geq 1} \frac{6}{m} a_m^\dagger a_m. \quad (9)$$

Sandwiching Eq.(4) with the Fock vacuum $|0\rangle$ and using $a_n|0\rangle = \langle 0|a_n^\dagger = 0$, we get

$$\sigma(\sigma^2 + g - \frac{6\zeta(2)\hbar^2}{g + 3\sigma^2}) = 0, \quad (10)$$

where $\zeta(2) = \frac{\pi^2}{6}$ for the Riemann zeta function $\zeta(k) \equiv \sum_{n \geq 1} \frac{1}{n^k}$. The second term in the expression of $\tilde{\alpha}_1(n)$ is derived from the last term of the constraint equation (4). Both the terms do not contribute to the vacuum expectation value of a_0 . So we neglect the terms in the following. Equation (10) has non-zero solutions

$$\sigma^2 = \frac{-2g + \sqrt{g^2 + 3\pi^2\hbar^2}}{3}, \quad (11)$$

and one finds a critical coupling $\lambda_{cr} = 24\mu^2/\hbar$ from $\sigma = 0$. In the limit $g \rightarrow -\infty$, we have $\sigma = \pm(\sqrt{-g} + \frac{3}{2}\zeta(2)\frac{\hbar^2}{\sqrt{-g^3}} + \dots)$. This asymptotic behavior agrees with Ref. [12] in which certain Feynman diagrams are calculated to the one-loop level within the framework of discretized light-front quantization. One also finds that $\sigma \sim (\lambda - \lambda_{cr})^\beta$ with $\beta = 1/2$ near the critical point. The same critical exponent has already been obtained in Ref.[12], but the value $\beta = 1/2$ disagrees with the expected value $\beta = 1/8$ indicated from the Ising model [15].

At $O(\hbar^2)$, similarly, one finds from the constraint equation (4)

$$\tilde{\alpha}_2(n) = -(\frac{6}{nA})^2 a_0 a_n + \frac{108}{n^2 A^3} a_0^3 a_n, \quad (12)$$

and σ satisfies

$$\sigma^2 + g - \frac{\pi^2 \hbar^2}{g + 3\sigma^2} + \zeta(3)\hbar(\frac{6\hbar}{g + 3\sigma^2})^2 \frac{g}{g + 3\sigma^2} = 0. \quad (13)$$

At $O(\hbar^2)$ one again finds $\sigma \sim (\lambda - \lambda_{cr})^{1/2}$ near the critical point.

It is not so easy to proceed to the higher order of $\hbar$, because of appearance of terms including $a_m^\dagger a_m$ and a_0^3 in $\tilde{\alpha}_p$. However, it is possible to set safely $\sigma^3 = 0$ at the critical point. So we concentrate on calculating a critical coupling λ_{cr} . We then decompose the $\tilde{\alpha}_p(n)$ into

$$\tilde{\alpha}_p(n) = \frac{\alpha_p(n)}{A^p} a_0 a_n + O(a_m^\dagger a_m) + O(a_0^3) \quad (n \geq 1), \quad (14)$$

with mere number $\alpha_p(n)$, and neglect the $O(a_0^3)$ terms in the following. The second term $O(a_m^\dagger a_m)$ contains terms like $\sum_m a_m^\dagger a_m a_0 a_n / A^{p+1}$ and $\sum_{m,l} a_m^\dagger a_m a_l^\dagger a_l a_0 a_n / A^{p+2}$ that vanish when sandwiched by $\langle 0|$ and $a_n^\dagger |0\rangle$. If $\alpha_p(n)$ are found, Eq.(4) yields

$$\begin{aligned} 0 &= \sigma^3 + g\sigma - \sum_{p,n \geq 1} \frac{\alpha_p(n)}{n} \langle 0| A^{-p} a_0 a_n a_n^\dagger |0\rangle \hbar^p + O(\sigma^3) \\ &= \sigma^3 + g\sigma - \sum_{p,n \geq 1} \frac{\alpha_p(n)}{n} g^{-p} \sigma \hbar^{p+1} + O(\sigma^3). \end{aligned} \quad (15)$$

Now, we calculate $\alpha_p(n)$. It is easy to calculate the commutation relation (6) except for the following term

$$\sum_{m \geq 1} \frac{1}{m} (a_m [a_0, a_n] a_m^\dagger + a_m^\dagger [a_0, a_n] a_m)$$

$$\begin{aligned}
&= \sum_{p \geq 1} \frac{1}{n} \frac{\alpha_p(n)}{A^p} a_0 a_n \hbar^{p+1} + \sum_{m, p \geq 1} \left(\frac{2}{m} a_m^\dagger a_m + \frac{\hbar}{m} \right) \frac{\alpha_p(n)}{A^p} a_0 a_n \hbar^p \\
&- \sum_{m, p \geq 1} \frac{1}{m} \left[\left[\frac{\alpha_p(n)}{A^p} a_0 a_n, a_m \right], a_m^\dagger \right] \hbar^p + O(a_m^\dagger a_m) + O(a_0^3). \tag{16}
\end{aligned}$$

Collecting $\hbar^p$ terms in Eq.(6), one obtains

$$\alpha_p(n) = -\frac{6}{n} \alpha_{p-1}(n) + [A^{p-1} \sum_{m \geq 1} \sum_{k \geq 1} \left(\frac{1}{m} \left[\left[\frac{\alpha_k(n)}{A^k} a_0 a_n, a_m \right], a_m^\dagger \right] \hbar^k \right)]_p, \quad 1 \leq p \leq 4 \tag{17}$$

where $\alpha_0(n) = -1$ and the bracket $[B]_p$ means extracting a coefficient C of $a_0 a_n \hbar^p$ from the operator $B = C a_0 a_n \hbar^p + \dots$. This recursion relation is true for $1 \leq p \leq 4$, since it is derived with $\tilde{\alpha}_1$ and $\tilde{\alpha}_2$ which have no $O(a_m^\dagger a_m)$ term. Using

$$\frac{1}{A} a_m = a_m \frac{1}{A - \frac{6}{m} \hbar} + O(a_0^2), \quad \frac{1}{A} a_m^\dagger = a_m^\dagger \frac{1}{A + \frac{6}{m} \hbar} + O(a_0^2), \tag{18}$$

one obtains from Eq.(17)

$$\begin{aligned}
\alpha_3(n) &= \frac{4}{3} \left(\frac{6}{n} \right)^3 + \frac{1}{3} \frac{6}{n} \sum_{m \geq 1} \left(\frac{6}{m} \right)^2, \\
\alpha_4(n) &= -\frac{7}{3} \left(\frac{6}{n} \right)^4 - \frac{5}{6} \left(\frac{6}{n} \right)^2 \sum_{m \geq 1} \left(\frac{6}{m} \right)^2 - \frac{1}{2} \frac{6}{n} \sum_{m \geq 1} \left(\frac{6}{m} \right)^3.
\end{aligned} \tag{19}$$

For $p \geq 5$, on the other hand, the $O(a_m^\dagger a_m)$ terms in $\tilde{\alpha}_i$ ($i \geq 3$) give some modifications to the recursion relation. In principle, the recursion relation is obtainable also for $p \geq 5$, but in practice the calculation is very complicated. So we consider the case of $p \leq 4$.

Setting $\sigma^3 = 0$ in Eq.(15), we obtain an equation for critical coupling g_{cr} , which is valid up to $O(\hbar^6)$,

$$\begin{aligned}
0 &= 6 - \zeta(2) \left(\frac{x}{\pi} \right)^2 + \zeta(3) \left(\frac{x}{\pi} \right)^3 - \left(\frac{4}{3} \zeta(4) + \frac{1}{3} \zeta(2)^2 \right) \left(\frac{x}{\pi} \right)^4 + \left(\frac{7}{3} \zeta(5) + \frac{4}{3} \zeta(3) \zeta(2) \right) \left(\frac{x}{\pi} \right)^5 \\
&= 36 - x^2(1 - 0.23261x + 0.14444x^2 - 0.09913x^3), \tag{20}
\end{aligned}$$

where $x = (6\pi\hbar)/g_{cr}$. We adopt the Pade approximations [16] to an alternating series $1 - c_1 x + c_2 x^2 - c_3 x^3 + \dots$. This series is approximated into $[M/N]$, which denotes a rational equation of polynomials of degrees M and N :

$$\begin{aligned}
[i/1] \quad & 1 - c_1 x + \dots + (-1)^i c_i \frac{x^i}{1 + \frac{c_{i+1}}{c_i} x}, \\
[1/2] \quad & \frac{1 + \left(\frac{c_3 - c_1 c_2}{c_2 - c_1^2} - c_1 \right) x}{1 + \frac{c_3 - c_1 c_2}{c_2 - c_1^2} x + \left(\frac{c_3 - c_1 c_2}{c_2 - c_1^2} c_1 - c_2 \right) x^2}.
\end{aligned} \tag{21}$$

Lower (upper) bounds of $\lambda_{cr} = (4\mu^2 x)/\hbar$ are obtained from cases of $M + N = \text{even (odd)}$,

$$\begin{aligned} [0/0] \quad \lambda_{cr} &= 24\mu^2/\hbar, & [1/1] \quad \lambda_{cr} &= 28.8\mu^2/\hbar, \\ [0/1] \quad \lambda_{cr} &= 46.0\mu^2/\hbar, & [2/1] \quad \lambda_{cr} &= 32.0\mu^2/\hbar, \\ [1/2] \quad \lambda_{cr} &= 31.1\mu^2/\hbar. \end{aligned} \quad (22)$$

Our result is $28.8\mu^2/\hbar \leq \lambda_{cr} \leq 31.1\mu^2/\hbar$, which is consistent with the equal-time result [8], $22\mu^2/\hbar \leq \lambda_{cr} \leq 55.5\mu^2/\hbar$.

So far the symmetric ordering has been assumed in the constraint equation. We now consider another possible operator ordering that does not break the hermitian nature. Under the ordering the constraint equation becomes

$$\begin{aligned} 0 &= a_0^3 + ga_0 + \sum_{n \geq 1} \frac{1}{n} ((a_0 a_n a_n^\dagger + a_0 a_n^\dagger a_n + a_n a_n^\dagger a_0 + a_n^\dagger a_n a_0)(1+s) + (a_n a_0 a_n^\dagger \\ &+ a_n^\dagger a_0 a_n)(1-2s) - 3\hbar a_0) + \sum_{k,l,m \neq 0} \frac{\delta_{k+l+m,0}}{\sqrt{|klm|}} a_k a_l a_m \equiv \Phi_s(a, a^\dagger) \end{aligned} \quad (23)$$

for a real parameter s . When $s = 0$, this ordering agrees with the symmetric one. Equations corresponding to (15) and (17) are

$$0 = \sigma^3 + g\sigma - \sum_{p,n \geq 1} \frac{\alpha_p(n)}{n} g^{-p} \sigma \hbar^{p+1} (1-2s) + O(\sigma^3), \quad (24)$$

$$\alpha_p(n) = -\frac{6}{n} \alpha_{p-1}(n) + (1-2s) [A^{p-1} \sum_{m \geq 1} \sum_{k \geq 1}^{p-2} (\frac{1}{m} [[\frac{\alpha_k(n)}{A^k} a_0 a_n, a_m], a_m^\dagger] \hbar^k)]_p, \quad (25)$$

for $1 \leq p \leq 4$. Since the double bracket term in Eq. (25) is $O(\hbar^3)$, α_1 and α_2 do not depend on s . At $O(\hbar)$ we have

$$\sigma(\sigma^2 + g - \frac{\pi^2 \hbar^2}{g + 3\sigma^2} (1-2s)) = 0, \quad (26)$$

so that $\lambda_{cr} = 24\mu^2/(\hbar\sqrt{1-2s})$ and $\beta = 1/2$ for $s < 1/2$. The spontaneous breaking does not occur for $s \geq 1/2$. When $-0.095 \leq s \leq 0.407$, the critical coupling calculated here reproduces the equal-time one $22\mu^2/\hbar \leq \lambda_{cr} \leq 55.5\mu^2/\hbar$. If the equal-time calculation yields λ_{cr} more accurately, one can find which s reproduces the λ_{cr} and can know whether

the symmetric ordering is the only ordering which reproduces the equal-time result. While λ_{cr} depends on s , β does not. This property persists for the higher order of $\hbar$. The present value of β is still different from the expected value $1/8$.

In this paper, we propose a systematic way of obtaining λ_{cr} and β within the framework of discretized light-front quantization. When the symmetric ordering is taken in the constraint equation, this formulation yields a reasonable value of λ_{cr} . The calculated λ_{cr} has an error much smaller than that of the equal-time calculation. This light-front formulation, however, does not reproduce the expected critical behavior $\sigma \sim (\lambda - \lambda_{cr})^{1/8}$. As long as the expansion (7) of $[a_0, a_n]$ is truncated at a finite order of $\hbar$, the non-zero solutions of σ obey an equation in which the higher order terms are added to Eq. (13). It is unlikely that in the equation the first term σ^2 is canceled by the higher order terms. So the unsatisfactory critical behavior $\sigma \sim (\lambda - \lambda_{cr})^{1/2}$ does not seem to break. This behavior is not changed by taking another operator ordering (23). This is an important problem to be solved. The first thing which we must do is to find the commutation relation $[a_0, a_n]$ without the truncation.

References

- [1] S.J. Chang, R.G. Root and T.M. Yan, Phys. Rev. **D7**, 1133(1973); S.J. Chang and T.M. Yan, Phys. Rev. **D7**, 1147(1973); T.M. Yan, Phys. Rev. **D7**, 1760(1973); T.M. Yan, Phys. Rev. **D7**, 1780(1973).
- [2] H.-C.Pauli and S.J.Brodsky, Phys.Rev. **D32**, 1993(1985); H.-C.Pauli and S.J.Brodsky, Phys.Rev. **D32**, 2001(1985); T. Eller, H.C. Pauli and S. Brodsky, Phys. Rev. **D35**, 1493(1987); T. Eller and H.C. Pauli, Z. Phys. **C42**, 59(1989); K. Hornbostel, S. Brodsky and H.Pauli, Phys. Rev. **D41**, 3814(1990). The heading is mislabeled in Table I: M/g should be replaced by $M^2/(g^2/\pi + m^2)$. This is informed by Dr. Hornbostel.

- [3] M. Burkardt, Nucl. Phys. **A504**, 762(1989).
- [4] K. Harada, T. Sugihara, M. Taniguchi and M. Yahiro, Phys. Rev. **D49**, 4226(1994);
T. Sugihara, M. Matsuzaki and M. Yahiro, Phys. Rev. **D50**, 5274(1994).
- [5] T. Maskawa and K. Yamawaki, Prog. Theor. Phys. **56**, 270(1976).
- [6] N. Nakanishi and K. Yamawaki, Nucl. Phys. **B122**, 15(1977); Y. Kim, S. Tsujimaru
and K. Yamawaki, Phys. Rev. Lett. **74**, 4771(1995).
- [7] K. Hornbostel, Phys. Rev. **D45**, 3781(1992).
- [8] S.J. Chang, Phys.Rev. **D13**, 2778(1976); J. Abad, J. G. Esteve and A. F. Pacheco,
Phys.Rev. **D32**, 2729(1985); M. Funke, U. Kaulfuss and H. Kummel, Phys.Rev. **D35**,
621(1987); H.Kroger, R. Girard, and G. Dufour,Phys.Rev. **D35**, 3944(1987).
- [9] T. Heinzl, S. Krusche, S. Simburger and E. Werner, Z. Phys.**C56**, 415(1992);
Phys.Lett.**B272**, 54(1991); Phys.Lett. **B275**, 410(1992); T. Heinzl, C. Stern, E.
Werner and B. Zeller mann, Z. Phys. **C72**, 353(1996).
- [10] A. Harindranath and J. P. Vary, Phys. Rev. **D36**, 1141(1987)1141; Phys. Rev.**D37**,
1076(1988); D. G. Robertson, Phys. Rev. **D47**, 2549(1993); M. Burkardt, Phys. Rev.
D47, 4628(1993).
- [11] C. M. Bender, S. S. Pinsky and B. van de Sande, Phys. Rev. **D48**, 816(1993); S. S.
Pinsky and B. van de Sande, Phys. Rev. **D49**, 2001(1994).
- [12] S.Pinsky, B. van de Sande and J. R. Hiller, Phys. Rev. **D51**, 726(1995).
- [13] X. Xu and H. J. Weber, Phys. Rev. **D52**, 4633(1995).
- [14] C. M. Bender, L. R. Mead and S. S. Pinsky, Phys. Rev. Lett. **56**, 2445(1986).
- [15] C. Domb and M. S. Green, *Phase Transitions and Critical Phenomena, Vol.6* (Aca-
demic, London, 1976).

[16] J. Killingbeck, *Microcomputer Quantum Mechanics* (Adam Hilger, Bristol, 1983).